

# MilkMatch™

## 20/20 AM BOV SAF DBZ

### CALF MILK REPLACER

#### MEDICATED

#### Dairy Herd and Beef Calf Milk Replacer

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves. Feed to provide 45.4 milligrams of lasalocid per 100 pounds (1 mg/2.2 lb) of body weight daily. Feed during periods of coccidiosis exposure or when experience indicates coccidiosis is likely to be a problem.

#### ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium) . . . . . 72.6 g/ton  
For preventing the development of house, stable, face, and horn flies in the manure of treated calves.

#### ACTIVE INGREDIENT

Diflubenuron (CAS #35367-38-5) . . . . . 13.6 g/ton

**WARNING:** Withdrawal period has not been established for lasalocid in pre-ruminating calves. Do not use in calves to be processed for veal.

- Do not feed to lactating cows.
- Do not feed to foals, dogs, and cats.

**CAUTION:** The safety of lasalocid in unapproved species has not been established.

#### GUARANTEED ANALYSIS

Crude Protein, not less than . . . . . 20.00%  
Crude Fat, not less than . . . . . 20.00%  
Crude Fiber, not more than . . . . . 0.15%  
Calcium, not less than . . . . . 0.75%  
Calcium, not more than . . . . . 1.25%  
Phosphorus, not less than . . . . . 0.70%  
Vitamin A, not less than . . . . . 35,000 IU/lb  
Vitamin D<sub>3</sub>, not less than . . . . . 7,500 IU/lb  
Vitamin E, not less than . . . . . 150 IU/lb

#### INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Animal and Vegetable Fat (Preserved with BHA and BHT), Dried Skimmed Milk, Lecithin, Dicalcium Phosphate, Calcium Carbonate, Yeast Extract, *Saccharomyces cerevisiae*, L-Lysine Monohydrochloride, DL-Methionine, L-Threonine, Vitamin A Supplement, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement, Ascorbic Acid, Magnesium Oxide, Zinc Sulfate, Ferrous Sulfate, Niacin Supplement, Manganese Sulfate, Calcium Pantothenate, Vitamin B<sub>12</sub> Supplement, Thiamine Mononitrate, Riboflavin Supplement, Biotin, Copper Sulfate, Pyridoxine Hydrochloride, Ethylenediamine Dihydrochloride, Folic Acid, Choline Chloride, Cobalt Sulfate, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Selenium Yeast, Sodium Aluminosilicate, Mono and Diglycerides of Edible Fats or Oils, Natural and Artificial Flavors.

#### SEE FEEDING DIRECTIONS ON REVERSE SIDE

Net Weight 25 lb (11.34 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

Manufactured for:

Augusta Cooperative Farm Bureau, Inc.  
Staunton, VA 24401  
MADE IN THE U.S.A.

502018

REV 0223

FI-015

## FEEDING PROGRAM

No whole milk is required after the colostrum feeding period when calf milk replacer is used.

#### MIXING AND FEEDING INSTRUCTIONS

Mix milk replacer at a rate of 12 oz of dry powder\* per 2 quarts of water (110° F).

Calves will vary in feed requirements depending on size and age. For proper medication levels (Lasalocid at 45.4 mg/ 100 lb body weight) - see chart. Each quart of milk replacer will contain 13.61 mg of lasalocid and 2.55 mg of diflubenuron.

Calf Weight (lb)	Quarts Fed per Feeding - Twice a Day
120	2

\*Dry powder should be weighed to determine recommended amount.

#### FOR BEST RESULTS:

- Provide starter (grain mix) for calves when they are about 4 days of age and feed free choice up to appetite.
- Calves can be weaned when they consume 2 to 3 pounds of starter feed for 3 consecutive days. Weaning time may need to be adjusted for smaller calves.

#### GOOD MANAGEMENT PRACTICES:

- Good newborn care is crucial to maximize calf survival rates. Immediate feeding of colostrum is one of the most important steps to increase the survival and health of newborn calves. Feed 2 quarts of high quality, first milking colostrum within the first 30 minutes after birth. Feed an additional 2 quarts at 6 to 12 hours of age. Continue feeding colostrum through the third day after birth.
- Begin your milk replacer program on the 4th day. Feed twice daily, at regular intervals, according to label directions. Wash equipment thoroughly after each usage.
- Have fresh, clean water available at all times.
- Start feeding product with ClariFly® early in the spring, 30 days before flies begin to appear, and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.
- In some cases, supplemental fly control measures may be needed in and around barns and calf hutches to control adult house and stable flies, which can breed in other decaying organic matter or silage in the area, or that migrate in from other locations.
- Do not feed hay until after weaning, when calves are consuming adequate amounts of dry starter.
- Calves should be housed in clean, dry individual pens that are well ventilated and draft free. Avoid sudden changes in temperature.

REV 0223

502018

ClariFly® is a registered trademark of Wellmark International.  
MilkMatch™ is a trademark of Augusta Cooperative Farm Bureau, Inc.