



Power-Lyx® 25% ANP Cattle Supplement

For Beef Cattle on Pasture

Guaranteed Analysis:

Crude Protein (Min)	25.00%
Crude Fat (Min)	4.00%
Crude Fiber (Max)	5.00%
Calcium (Min)	0.50%
Calcium (Max)	1.00%
Phosphorus (Min)	0.50%
Salt (Min)	2.50%
Salt (Max)	3.50%
Magnesium (Min)	3.00%
Potassium (Min)	0.50%
Copper (Min)	30 ppm
Manganese (Min)	100 ppm
Selenium (Min)	3.3 ppm
Zinc (Min)	75 ppm
Vitamin A (Min)	25,000 IU/lb
Vitamin D-3 (Min)	4,000 IU/lb
Vitamin E (Min)	4 IU/lb

Power-Lyx® is a registered trademark of Ridley Block Operations.

Manufactured For:
Augusta Cooperative Farm Bureau, Inc.
Staunton, VA 24401

Ingredients:

Molasses Products, Processed Grain By-Products, Animal Protein Products, Magnesium Oxide, Salt, Hydrolyzed Vegetable Oil, Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Roughage Products, Sodium Selenite, Manganese Sulfate, Copper Sulfate, Zinc Sulfate, Vitamin A Supplement, Manganous Oxide, Vitamin D-3 Supplement, Zinc Oxide, Vitamin E Supplement, Mineral Oil, Ethylenediamine Dihydroiodide, Cobalt Carbonate, and Calcium Iodate.

Introductory Period: Following a 7 to 10 day acclimation period, average consumption should be 1 to 2 pounds for mature cattle with unrestricted forage supplies. Position the blocks 50 to 75 feet from loafing, feeding and watering areas. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head daily.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

Feeding Recommendations:

1. Feed 1 block per 15 to 30 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Cattle should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle should consume 1 to 2 pounds per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate free choice mineral supplement in addition to this supplement.

For additional information, please contact your Augusta Co-op representative.

Net Weight: 200 LBS. (90.7 KG)

56620