

INTERVET INC., MERCK ANIMAL HEALTH
556 MORRIS AVE., SUMMIT, NJ, 07901

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 908-473-5560

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

VISION[®] 7 WITH SPUR^{®*}
Intervet/Merck Animal Health

Cattle Vaccine

CLOSTRIDIUM CHAUVOEI-SEPTICUM-NOVYI-SORDELLII-PERFRINGENS TYPES C & D BACTERIN-TOXOID

For use in healthy cattle and sheep as an aid in preventing disease caused by *Clostridium chauvoei* (Blackleg), *septicum* (Malignant edema), *novyi* (Black disease), *sordellii* and *perfringens* Types C & D (Enterotoxemia).

DOSE: Cattle-2mL, Sheep-2mL. Inject subcutaneously. Repeat in 3 to 4 weeks. Annual revaccination is recommended. Shake well before using. Store at 35° to 45°F (2° to 7°C). Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

ANTIDOTE: Epinephrine. Contains formaldehyde as preservative.

*Adjuvant-Intervet's Proprietary Technology.

FOR ANIMAL USE ONLY

Intervet Inc., Omaha, NE 68103 USA

U.S. Vet Lic. No. 165A

1-800-521-5767

		Code	
10 Doses	20 mL	069130	06000958-06
50 Doses	100 mL	069131	06010958-05
250 Doses	500 mL	069132	06020958-05

NAC No.: 1047407.2