

WARNING: Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children.
Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information or to obtain MSDS, contact Durvet at 1-800-821-5570.

PRECAUTIONS: Ivermectin Paste (ivermectin) has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.
Environmental Safety: Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

INFORMATION FOR HORSE OWNERS: Swelling and itching reactions after treatment with Ivermectin Paste (ivermectin) have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp. microfilariae). These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions

occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with Ivermectin Paste (ivermectin). Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

Apple
Flavored!



IVERMECTIN (ivermectin) PASTE 1.87%

Net Wt. 0.21 oz (6.08 g)

ANTHELMINTIC & BOTICIDE

NDC 30798-835-81

Removes worms and bots
with a single dose.

Contents will treat up to
1250 lb body weight

For Oral Use In Horses Only
ANADA# 200-326, Approved by FDA



IVERMECTIN
(ivermectin) PASTE 1.87%

IVERMECTIN (ivermectin) PASTE 1.87%

ANTHELMINTIC & BOTICIDE



Distributed by: DURVET, INC.
Blue Springs, MO 64014

8IVE015

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Ivermectin Paste (ivermectin) provides effective treatment and control of the following parasites in horses. **Large Strongyles** (adults)- *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus*; **Small Strongyles** (adults, including those resistant to some benzimidazole class compounds)- *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*; *Cylicocyclus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus*, and *C. brevicapsulatus*; *Cylicodontophorus* spp.; *Cylicostephanus* spp. including *C. callicatus*, *C. goldi*, *C. longibursatus*, and *C. minutus*; **Small Strongyles** - Fourth stage larvae; **Pinworms** (adults and fourth-stage larvae)-*Oxyuris equi*; **Ascarids** (adults, third-stage and fourth-stage larvae)-*Parascaris equorum*; **Hairworms** (adults)-*Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults)-*Habronema muscae*; **Bots** (oral and gastric stages)- *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and fourth stage larvae)- *Dictyocaulus arnfieldi*; **Intestinal Threadworms** (adults)-*Strongyloides westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by **neck threadworm** microfilariae, *Onchocerca* sp.

DOSAGE AND ADMINISTRATION: This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight

marking on the syringe plunger delivers enough paste to treat 250 lb body weight.

(1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. (2) Lock the ring in place by making a 1/4 turn to the right. (3) Make sure that the horse's mouth contains no feed. (4) Remove the cover from the tip of the syringe. (5) Insert the syringe tip into the horse's mouth at the space between the teeth. (6) Depress the plunger as far as it will go, depositing paste on the back of the tongue. (7) Immediately raise the horse's head for a few seconds after dosing.

PARASITE CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. Ivermectin Paste (ivermectin) effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

PRODUCT ADVANTAGES: **Broad-spectrum Control** - Ivermectin Paste (ivermectin) kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose. Ivermectin Paste (ivermectin) is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

ANIMAL SAFETY: Ivermectin Paste (ivermectin) may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

April, 2006

Manufactured by:
Bimeda-MTC Animal Health Inc.
Cambridge, Ontario, Canada N3C 2W4

Bimeda-MTC Animal Health Inc.
is a Division of Cross Vetpharm Group Ltd.

UPC CODE
7 45801 11048 9

IVERMECTIN

(ivermectin) PASTE 1.87%

ANTHELMINTIC & BOTICIDE

For Treatment of Large Strongyles, Small Strongyles, Pinworms, Roundworms (Ascarids), Hairworms, Neck Threadworms, Large-mouth Stomach Worms and Bots. See carton or package insert for complete indications and use directions. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

WARNING: Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children.

Refrain from smoking and eating when handling.
Wash hands after use. Avoid contact with eyes.

NDC 30798-835-81



www.durvet.com

ANADA# 200-326, Approved by FDA
For Oral Use in Horses Only

Net Weight: 0.21 oz (6.08 g)

Manufactured by: Bimeda-MTC Animal Health Inc.
Cambridge, Ontario, Canada N3C 2W4

Bimeda-MTC Animal Health Inc. is a Division of Cross Vetpharm Group Ltd.

Distributed by: DURVET, INC.
Blue Springs, MO 64014

8IVE014

Iss.04.06



IVERMECTIN PASTE 1.87% (ivermectin)

ANTHELMINTIC & BOTICIDE

Removes worms and bots with a single dose.

Contents will treat up to 1250 lb body weight

For Oral Use In Horses Only

ANADA# 200-326, Approved by FDA

Net Wt. 0.21 oz (6.08 g)

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Ivermectin Paste 1.87% (ivermectin) provides effective treatment and control of the following parasites in horses.

Large Strongyles (adults)- *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus*; **Small Strongyles** (adults, including those resistant to some benzimidazole class compounds)- *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*; *Cylicocycylus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus*, and *C. brevicapsulatus*; *Cylicodontophorus* spp.; *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus*, and *C. minutus*; **Small Strongyles** - Fourth stage larvae; **Pinworms** (adults and fourth-stage larvae)-*Oxyuris equi*; **Ascarids** (adults, third-stage and fourth-stage larvae)-*Parascaris equorum*; **Hairworms** (adults)-*Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults)-*Habronema muscae*; **Bots** (oral and gastric stages)- *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and fourth stage larvae)- *Dictyocaulus arnfieldi*; **Intestinal Threadworms** (adults)-*Strongyloides westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by **neck threadworm** microfilariae, *Onchocerca* sp.

DOSAGE AND ADMINISTRATION: This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight.

(1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. (2) Lock the ring in place by making a 1/4 turn to the right. (3) Make sure that the horse's mouth contains no feed. (4) Remove the cover from the tip of the syringe. (5) Insert the syringe tip into the horse's mouth at the space between the teeth. (6) Depress the plunger as far as it will go, depositing paste on the back of the tongue. (7) Immediately raise the horse's head for a few seconds after dosing.

PARASITE CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. Ivermectin Paste 1.87% (ivermectin) effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

PRODUCT ADVANTAGES: Broad-spectrum Control - Ivermectin Paste 1.87% (ivermectin) kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose. Ivermectin Paste 1.87% (ivermectin) is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

ANIMAL SAFETY: Ivermectin Paste 1.87% (ivermectin) may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

WARNING: Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children.

Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes.

The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information or to obtain a MSDS, contact Durvet at 1-800-821-5570.

PRECAUTIONS: Ivermectin Paste 1.87% (ivermectin) has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

Environmental Safety: Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

INFORMATION FOR HORSE OWNERS: Swelling and itching reactions after treatment with Ivermectin Paste 1.87% (ivermectin) have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp. microfilariae). These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with Ivermectin Paste 1.87% (ivermectin). Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

Manufactured by:
Bimeda-MTC Animal Health Inc.
Cambridge, Ontario, Canada N3C 2W4

Bimeda-MTC Animal Health Inc.
is a Division of Cross Vetpharm Group Ltd.

8IVE017

Iss.04.06



Distributed by: DURVET, INC.
Blue Springs, MO 64014