

INTERVET INC., MERCK ANIMAL HEALTH

556 MORRIS AVE., SUMMIT, NJ, 07901

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 908-473-5560

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

REVALOR®-G

Intervet/Merck Animal Health

(trenbolone acetate and estradiol)

NADA # 140-897, Approved by FDA

For Pasture Cattle (Slaughter, Stocker & Feeder Steers and Heifers)

DESCRIPTION

Revalor®-G (trenbolone acetate and estradiol) is an implant containing 40 mg of trenbolone acetate and 8 mg estradiol. Each implant consists of 2 small yellow pellets. Ten implants are provided in a cartridge.

Manufactured by a non-sterilizing process.

Restricted Drug (California) - use only as directed.

INDICATIONS FOR USE

Increases rate of weight gain in pasture cattle (slaughter, stocker, and feeder steers and heifers) in a slow-release delivery system.

Do not use in veal calves. Effectiveness and animal safety in veal calves have not been established.

DOSAGE

One implant containing 40 mg trenbolone acetate and 8 mg estradiol is administered to each animal. The 2 pellets which make up the dosage of Revalor®-G are contained in one division of the multiple dose cartridge. Ten doses are in each cartridge. The cartridge is designed to be used with a special implant gun.

ROUTE OF ADMINISTRATION:

The implant is placed under the skin on the posterior aspect of the ear by means of a special implanter available from Intervet Inc. With the animal suitably restrained, the skin on the outer surface of the ear should be cleaned. The implant is then administered by the method shown in the diagram below.




Fig. 1 - Ear of Bovine Ready for Implantation

SITE OF IMPLANTATION

After appropriately restraining the animal to allow access to the ear, cleanse the skin at the implant needle puncture site. It is subcutaneous between the skin and cartilage on the back side of the ear and below the midline of the ear. The implant must not be placed closer to the head than the edge

of the cartilage ring farthest from the head. The location of insertion of the needle is a point toward the tip of the ear and at least a needle length away from the intended deposition site. Care should be taken to avoid injuring the major blood vessels or cartilage of the ear.



Fig. 2 - Rear View of the Bovine Ear Showing the Site for Insertion of the Implanter Needle.

METHOD OF USE

1. Do not remove the cap of the cartridge containing the implants.
2. Place the cartridge (D) with the capped end to the front into slot at the top of the implanter magazine marked (A) on the diagram.
3. Gently push the cartridge into the slot until it clicks into place.
4. The implanter is then ready for use.
5. Take the ear of the animal firmly with the free hand in the manner shown in Fig. 1. Then insert the needle into the subcutaneous tissue at the point indicated in Fig. 2.
6. After inserting the needle to its full extent, squeeze the trigger (E) gradually. Allow the pellets of the implant to be deposited in a single row.
7. Withdraw the implanter. This will advance the cartridge one groove in the magazine and the next implant is now ready for use.
8. When all the implants have been administered, the cartridge will discharge out the bottom of the magazine and may be replaced by a new one.
9. To change the needle, loosen the needle locking nut labeled (F) in Fig. 3 and replace the needle. Tighten the nut finger tight and the implanter is ready for use.

A - Magazine aperture
B - Needle
C - Cartridge cap
D - Cartridge
E - Trigger
F - Needle locking nut




Fig. 3 - Diagram of the Implanter and Cartridge

WARNING

Not to be used in animals intended for subsequent breeding, or in dairy animals. For Animal Treatment Only. Not for Use in Humans.

Implant pellets in the ear only. Any other location is in violation of Federal Law. Do not attempt salvage of implanted site for human or animal food.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

STORAGE CONDITIONS

Store unopened product at or below 25°C (77°F). Avoid excessive heat and humidity. Use product before the expiration date printed on the label and on the cartridge pouch. Opened cartridges may be stored in the foil pouch protected from light in the refrigerator (2-8°C/36-47°F) for up to 6 months.

HOW SUPPLIED

Available as a box of 10 x 10 cartridge implants.

Revalor is a registered trademark of Intervet Inc. or an affiliate.

Made in Austria by: Intervet GesmbH

Distributed by: Intervet Inc., Millsboro, DE 19966

481200-E, 781200-D 08.06

009832 INW240 01

NAC No.: 1047379.1