



1906BG SAFEGUARD 0.5% CATTLE

MEDICATED

Swine/Beef/Dairy/Horse Dewormer

For the removal and control of lungworms (*Dictyocaulus viviparus*); barberpole worms (*Haemonchus contortus*); brown stomach worms (*Ostertagia ostertagi*); small stomach worms (*Trichostrongylus axei*); hookworms (*Bunostomum phlebotomum*); thread-necked intestinal worms (*Nematodirus helvetianus*); small intestinal worms (*Cooperia punctatata* and *Cooperia oncophora*); bankrupt worms (*Trichostrongylus colubriformis*); and nodular worms (*Oesophagostomum radiatum*).

ACTIVE DRUG INGREDIENT

Fenbendazole 4635 g/ton

Guaranteed Analysis:

Crude Protein	(Min)	13.0 %
Crude Fat	(Min)	3.7 %
Crude Fiber	(Max)	8.5 %
Calcium(Ca)	(Min) 4.6	(Max) 5.3 %
Phosphorus(P)	(Min)	0.56 %
Potassium(K)	(Min)	0.73 %

Ingredients:

Processed Grain By-Products, Calcium Carbonate, SAFEGUARD 20% and Vegetable and Animal Fat (Preserved with BHT).

Directions for use in Beef and Dairy Cattle

Feed at the rate of 0.10 pound per 100 pounds of bodyweight.

LIMITATIONS: Feed as sole ration or topdress for one day. Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle, the milk discard time is zero hours. A withdrawal time has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

NOTE: Consult veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Directions for use for Swine

For Growing Pigs, Gilts, Pregnant Sows, and Boars - 9 mg active drug ingredient over a Fenbendazole per kg body weight (4.08 mg Fenbendazole per pound) to be fed as the sole ration period of 3 to 12 days. Example - 200 lb pig consuming 8 lbs of feed should receive 1.92 ounces of SafeGuard 0.5% each day for 3 days. No prior withdrawal of feed or water necessary.

LIMITATIONS: Residue Warning - There is no pre-slaughter withdrawal period as SafeGuard 0.5% can be fed to day of slaughter.

Directions for use for Equines

5 mg active ingredient Fenbendazole per kg body weight (2.27 mg Fenbendazole per pound) in a one (1) day treatment for large strongyles, small strongyles and pinworms. Example: a 1000 lb horse should receive 1.0 pounds of SafeGuard 0.5% in a one day treatment. For ascarids the recommended dose is 10 mg Fenbendazole per kg body weight (4.54 mg Fenbendazole per pound) in a one day treatment. Example: a 1000 lb horse should receive 2 pounds of SafeGuard 0.5% in a one day treatment. All horses must be eating normally to ensure that each animal consumes an adequate amount of medicated feed. Regular deworming at intervals of 6 to 8 weeks may be required due to the possibility of reinfection. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

LIMITATIONS: Do not use in horses intended for food.

MANUFACTURED FOR:
IVESCO, LLC
IOWA FALLS, IA 50126

NET WT 25 LB (11.36 KG)

MEDICATED