



A Liquid Topical Flea and Lice Product for Dogs

READ THE ENTIRE LABEL BEFORE EACH USE
ACTIVE INGREDIENTS:

Imidacloprid:	9.10%
Pyriproxyfen:	0.46%
OTHER INGREDIENTS:	90.44%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
HAZARDS TO DOMESTIC ANIMALS. For external use only. Do not use on puppies under 7 weeks of age or weighing less than 3 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing dogs. Individual sensitivities, while rare, may occur after using ANY pesticide products for Dogs. If signs persist, or become more severe, consult a veterinarian immediately. If your dog is on medication, consult your veterinarian before using this or any other product. **RESTRICTIONS:** Use only on dogs 7 weeks and older weighing over 3 lbs. Do not use on other animals. Do not exceed labeled dosage amount for dogs. Do not apply more than one tube per treatment. Do not have contact or allow children to have contact with treated area until completely dry. Do not treat more than 1 dog per tube.

FIRST AID	
If Swallowed:	Call a poison control center or a doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If In Eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If On Skin:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies call 1-844-685-9173.	
NOTE TO PHYSICIAN	
Treat the patient symptomatically.	

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **DO NOT ALLOW CHILDREN TO APPLY PRODUCT.** To prevent harm to you and your dog, read the entire label and package insert before each use. Follow all directions and precautionary statements carefully. Use **ONLY** on Dogs and Puppies 7 Weeks and Older and Weighing Over 3 lbs.

Side Effects: Monitor your dog after application. Side effects, although very rare, may include signs of skin irritations such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-844-685-9173.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. **Pesticide storage:** Store in a cool, dry place inaccessible to children and pets. **Pesticide Disposal and Container Handling:** Nonrefillable container. **If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian immediately.

If you have questions or comments about this product, please write:
Promika, LLC
1204 Village Market PI #273, Morrisville, NC 27560

CONTAINS IMIDACLOPRID AND PYRIPROXYFEN, THE ACTIVE INGREDIENTS USED IN ADVANTAGE® II FOR DOGS

SMALL For Use **ONLY** on Dogs and Puppies 7 Weeks and Older Weighing **3 to 10 lbs.**

4-Pack of 0.014 fl oz
Net Contents: 0.056 fl oz

MEDIUM For Use **ONLY** on Dogs and Puppies 7 Weeks and Older Weighing **11 to 20 lbs.**

4-Pack of 0.034 fl oz
Net Contents: 0.136 fl oz

LARGE For Use **ONLY** on Dogs and Puppies 7 Weeks and Older Weighing **21 to 55 lbs.**

4-Pack of 0.084 fl oz
Net Contents: 0.336 fl oz

X-LARGE For Use **ONLY** on Dogs and Puppies 7 Weeks and Older Weighing **Over 55 lbs.**

4-Pack of 0.135 fl oz
Net Contents: 0.54 fl oz



FLEAS



EGGS



LARVAE



LICE

DIRECTIONS FOR USE | 3-20 LBS

Separate paper from corner of blister package. Peel back paper and take out the tube. Remove one applicator tube from the package. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube. Turn the cap around and place other end of cap back on tube. Twist cap to break seal, then remove cap from tube. The dog should be standing for easy application. Part the hair on the dog's back, between the shoulder blades, until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. Do not get this product in your dog's eyes, or allow your dog to ingest this product. Do not allow the product to run off. Discard empty tube as described in the Storage and Disposal section. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not re-treat more often than once every seven days. After flea control is attained, return to a monthly retreatment schedule.

DIRECTIONS FOR USE | 21+ LBS

Separate paper from corner of blister package. Peel back paper and take out the tube. Remove one applicator tube from the package. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube. Turn the cap around and place other end of cap back on tube. Twist cap to break seal, then remove cap from tube. The dog should be standing for easy application. Apply the entire contents of the tube evenly to three or four spots on the top of the back from the shoulder to the base of the tail. At each spot, part the hair until the skin is visible. Place the tip of the tube on the skin and gently squeeze to expel a portion of the solution on the skin. Do not apply any excessive amount of solution at any one spot that could cause some of the solution to run off the side of the dog. Do not get this product in your dog's eyes, or allow your dog to ingest this product. Discard empty tube as described in the Storage and Disposal section. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. Do not re-treat more often than once every seven days. After flea control is attained, return to a monthly retreatment schedule.



In Case of Emergency,
Call: 1-844-685-9173